Information sheet for customer

Tsumura-Kampo Shosaikoto[小柴胡湯] Extract Granules

[Category 2 OTC medicines]

INDICATIONS

Following symptoms in persons with moderate physical strength who complaint pain and discomfort in the solar plexus and flank, and suffered from loss of appetite, bitter sensation in the mouth and furry white tongue:

Loss of appetite, nausea, gastritis, gastric pain, gastrointestinal weakness, fatigue, and late stage of common cold.



! CAUTION FOR USE

Who should seek consultation

- 1. The following persons should consult a physician, pharmacist, or registered distributor before taking this medicine.
 - (1) Persons undergoing medical treatment from a physician.
 - (2) Pregnant women or women suspected of being pregnant.
 - (3) Persons who is physically weak (declining of strength or constitutional weakness).
 - (4) Elderly persons.
 - (5) Persons who have experienced rash, redness, itching, etc. due to medicines or something.
 - (6) Persons with the following symptoms: edema.
 - (7) Persons diagnosed as having the following: liver disease, hypertension, heart disease, kidney disease.
 - (8) Person who is treated with the interferon preparation.
- 2. The following symptoms may be adverse reactions to this medication. If any of these symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist, or registered distributor. Take the instruction leaflet with you.

Related areas	Symptoms
Skin	Rash, redness, itching
Others	Frequent urination, urination pain, bloody urine, feeling of
	residual urine

In very rare cases the following serious symptoms may occur. In this case, consult a physician immediately.

iniculatory.		
Symptoms' name	Symptoms	
Interstitial pneumonia	When go upstairs or exert yourself too much, symptoms such	
	as shortness of breath, breathing difficulties, dry cough, and	
	fever will appear. They will appear suddenly and persist.	
Pseudoaldosteronism,	In addition to listlessness, stretched feeling, and stiffness of	
myopathy	hands and feet, feeling of weakness and muscular pain will	
	appear and gradually worsen.	
Hepatic function failure	Following symptoms appear: fever, itching, rash, jaundice	
	(yellowing of skin and white of eyes), brown urine, whole-body	
	tiredness, loss of appetite.	

- 3. Stop taking this medicine and consult a physician, pharmacist or registered distributor if symptoms do not improve after taking this medicine for about 1 month (1 week if you take this medicine for late phase of common cold). Take this leaflet with you.
- 4. If you take this medicine regularly for a long periods, consult a physician, pharmacist or registered distributor.

DOSAGE

Take the following dosages with water or warm water before a meal.

Age	One dosage	Daily dosage
15 years and over	1 packet(1.875 g)	
7-14 years	2/3 packet	2 times a day
4-6 years	1/2 packet	
2-3 years	1/3 packet	
Less than 2 years	Do not use	

[Cautions] This medication should be given to children only under adult supervision

INGREDIENTS

2 packets (3.75 g) of this medicine contain 2.25 g of Shosaikoto extract of the following mixed crude drugs;

JP Bupleurum Root	3.5 g
JP Pinellia Tuber	2.5 g
JP Scutellaria Root	1.5 g
JP Jujube	1.5 g
JP Ginseng	1.5 g
JP Glycyrrhiza	1.0 g
JP Ginger	0.5 g

(JP: The Japanese Pharmacopoeia)

Inactive ingredients: JP Magnesium Stearate, JP Lactose Hydrate, Sucrose Esters of Fatty Acids

PRECAUTIONS FOR STRAGE AND HANDLING

- (1) Store in a cool, dry place free from direct sunlight.
- (2) Keep all medication out of the reach of children.
- (3) When you take this medicine in divided dose, fold over the opened edge of the packet and store it. The remaining medicine should be taken within 2 days.
- (4) Since the medicine contains crude drugs, the granules color may vary for individual products, however this has no influence on the efficacy of this medicine.
- (5) Do not take this medicine after the expiration date.

Production of this product has been approved for domestic sales in accordance with the Pharmaceutical Affairs Law of Japan.

