

Information sheet for customer

Tsumura-Kampo Hachimijioganryo[八味地黄丸料] Extract Granules A

Tsumura-Kampo Keate- Granules [ケアテ顆粒]

【Category 2 OTC medicines】

INDICATIONS

Following symptoms in persons with moderate or poor physical strength who tend to get tired easily, tend to be sensitive to cold in hands and feet, have decreased or increased urine volume, and/or sometimes have dry mouth: Leg pain, lower back pain, numbness, blurred eye in elderly, itching, difficulty in urination, residual urine, nocturia, frequent urination, edema, associated symptoms of hypertension (stiff shoulder, heavy head, ear noise), and mild incontinence.



! CAUTION FOR USE

●To be avoided

Failure to observe the following may result in the symptoms worsening or the occurrence of adverse reactions/accidents.

- (1) Persons with weak stomach and intestine.
- (2) Persons who have a tendency to diarrhea.

●Who should seek consultation

1. The following persons should consult a physician, pharmacist, or registered distributor before taking this medicine.
 - (1) Persons undergoing medical treatment from a physician.
 - (2) Pregnant women or women suspected of being pregnant.
 - (3) Red faced persons with hot flashes who have fulfilling physical performance.
 - (4) Persons who have experienced rash, redness, itching, etc. due to medicines or something.
2. The following symptoms may be adverse reactions to this medication. If any of these symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist, or registered distributor of this medication. Take the instruction leaflet with you.

Related areas	Symptoms
Skin	Rash, redness, itching
Gastrointestinal system	Loss of appetite, gastric distress, abdominal pain
Others	Palpitations, rush of blood to the head, cheilitis, numbness of the tongue

3. A following symptom may appear after taking this medicine. If the symptom continues or worsen, consult a physician, pharmacist or registered distributor. Take this leaflet with you: diarrhea
4. Stop taking this medicine and consult a physician, pharmacist or registered distributor if symptoms do not improve after taking this medicine for about 1 month. Take this leaflet with you.

DOSAGE

Take the following dosages with water or warm water before a meal.

Age	One dosage	Daily dosage
15 years and over	1 packet (1.875 g)	2 times a day
7-14 years	2/3 packet	
4-6 years	1/2 packet	
2-3 years	1/3 packet	
Less than 2 years	Do not use	

[Cautions] This medication should be given to children only under adult supervision

INGREDIENTS

2 packets (3.75 g) of this medicine contain 2.0 g of Hachimijiogan/Keate extract of the following mixed crude drugs;

JP Rehmannia Root	3.0 g
JP Cornus Fruit	1.5 g
JP Dioscorea Rhizome	1.5 g
JP Alisma Rhizome	1.5 g
JP Poria Sclerotium	1.5 g
JP Moutan Bark	1.25 g
JP Cinnamon Bark	0.5 g
JP Powdered Processed Aconite Root	0.25 g

(JP: The Japanese Pharmacopoeia)

Inactive ingredients: JP Magnesium Stearate, JP Lactose Hydrate, Sucrose Esters of Fatty Acids.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store in a cool, dry place free from direct sunlight.
- (2) Keep all medication out of the reach of children.
- (3) When you take this medicine in divided dose, fold over the opened edge of the packet and store it.
The remaining medicine should be taken within 2 days.
- (4) Since the medicine contains crude drugs, the granules color may vary for individual products, however this has no influence on the efficacy of this medicine.
- (5) Do not take this medicine after the expiration date.

Production of this product has been approved for domestic sales in accordance with the Pharmaceutical Affairs Law of Japan.



Tsumura&Co. 2-17-11, Akasaka, Minato-ku, Tokyo107-8521, JAPAN

<https://www.tsumura.co.jp/>