

Information sheet for customer

Tsumura-Kampo-Liquid Kakkonto 2 [葛根湯 2]

【Category 2 OTC medicines】

INDICATIONS

Following symptoms in person with moderate or more physical strength:

Initial stage of common cold(without spontaneous sweating), coryza, rhinitis, headache, shoulder stiffness, muscle pain, and pain in hands and shoulder.



! CAUTION FOR USE

●Who should seek consultation

1. The following persons should consult a physician, pharmacist, or registered distributor before taking this medicine.
 - (1) Persons undergoing medical treatment from a physician.
 - (2) Pregnant women or women suspected of being pregnant.
 - (3) Persons who is physically weak (declining of strength or constitutional weakness)
 - (4) Persons who have gastrointestinal weakness
 - (5) Persons with excessive sweating
 - (6) Elderly persons
 - (7) Persons who have experienced rash, redness, itching, etc. due to medicines or something.
 - (8) Persons with the following symptoms: edema, difficulty in urination.
 - (9) Persons diagnosed as having the following: hypertension, heart disease, kidney disease, thyroid disease.
2. The following symptoms may be adverse reactions to this medication. If any of these symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist, or registered distributor. Take the instruction leaflet with you.

Related areas	Symptoms
Skin	Rash, redness, itching
Gastrointestinal system	Nausea, loss of appetite, gastric distress

In very rare cases the following serious symptoms may occur. In this case, consult a physician immediately.

Symptoms' name	Symptoms
Pseudoaldosteronism, myopathy	In addition to listlessness, stretched feeling, and stiffness of hands and feet, feeling of weakness and muscular pain will appear and gradually worsen.
Hepatic function failure	Following symptoms appear: fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, whole-body tiredness, loss of appetite.

3. Stop taking this medicine and consult a physician, pharmacist or registered distributor if symptoms do not improve after taking this medicine for about 1 month (5 to 6 times if you take this medicine for the initial stage of the common cold, coryza, and headache). Take this leaflet with you.
4. If you take this medicine regularly for a long periods, consult a physician, pharmacist or registered distributor.

DOSAGE

Take the following dosages between meals.

Age	One dosage	Daily dosage
15 years and over	1 bottle	2 times a day
Less than 15 years	Do not use	

Shake the bottle before taking this medicine.

[Caution] Drink a whole bottle of this medicine at one time not leaving a part of it.

INGREDIENTS

Daily dosage 90 mL (45ml × 2 bottles) of this medicine contain 8.3 g of Kakkonto extract of the following mixed crude drugs;

JP Pueraria Root	8.0 g
JP Ephedra Herb	4.0 g
JP Jujube	4.0 g
JP Cinnamon Bark	3.0 g
JP Peony Root	3.0 g
JP Glycyrrhiza	2.0 g
JP Ginger	1.0 g

(JP: The Japanese Pharmacopoeia)

Inactive ingredients: Invert Type Liquid Sugar (Sucrose, Fructose, Glucose), JP Honey, JP Sodium Benzoate, JP Butyl Parahydroxybenzoate, Ethanol, Polyoxyethylene Hardened Castor Oil, JP Citric Acid Hydrate, JP Sodium Bicarbonate, Flavor, Ethyl Vanillin, Glycerin, Vanillin, Propylene Glycol.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store in a cool, dry place free from direct sunlight.
- (2) Keep all medication out of the reach of children.
- (3) Do not transfer this medicine to a different container (It may be misused or the quality may deteriorate).
- (4) Do not take this medicine after the expiration date.

Production of this product has been approved for domestic sales in accordance with the Pharmaceutical Affairs Law of Japan.

Sales agency:

Tsumura&Co.

2-17-11, Akasaka, Minato-ku, Tokyo 107-8521, JAPAN.

<http://www.tsumura.co.jp/>

Manufacturing and selling agency:

DAIDO PHARMACEUTICAL CORPORATION

214-1, Shimura, Katsuragi-Shi, Nara 639-2121, JAPAN.