

## Information sheet for customer

# Tsumura-Kampo Hangekobokuto[半夏厚朴湯] Extract Granules

【Category 2 OTC medicines】

### INDICATIONS

Following symptoms in person with moderate physical strength who feel depressed, experience feeling of a foreign body in the throat and esophagus, and sometimes experience palpitations, dizziness, nausea, etc:

Choking feeling in the throat, nervous gastritis, anxiety, cough, hoarseness, and morning sickness.



### ! CAUTION FOR USE

#### ●Who should seek consultation

1. The following persons should consult a physician, pharmacist, or registered distributor before taking this medicine.
  - (1) Persons undergoing medical treatment from a physician.
  - (2) Persons who have experienced rash, redness, itching, etc. due to medicines or something.
2. The following symptoms may be adverse reactions to this medication. If any of these symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist, or registered distributor. Take the instruction leaflet with you.

Related areas	Symptoms
Skin	Rash, redness, itching

3. Stop taking this medicine and consult a physician, pharmacist or registered distributor if symptoms do not improve after taking this medicine for about 1 month ( 5 to 6 days if you take this medicine for morning sickness). Take this leaflet with you.

### DOSAGE

Take the following dosages with water or warm water before a meal.

Age	One dosage	Daily dosage
15 years and over	1 packet(1.875 g)	2 times a day
7-14 years	2/3 packet	
4-6 years	1/2 packet	
2-3 years	1/3 packet	
Less than 2 years	Do not use	

[Cautions] This medication should be given to children only under adult supervision

## INGREDIENTS

2 packets (5.0 g) of this medicine contain 1.25 g of Hangekobokuto extract of the following mixed crude drugs;

JP Pinellia Tuber	3.0 g
JP Poria Sclerotium	2.5 g
JP Magnolia Bark	1.5 g
JP Perilla Herb	1.0 g
JP Ginger	0.5 g

(JP: The Japanese Pharmacopoeia)

Inactive ingredients: JP Magnesium Stearate, JP Lactose Hydrate, Sucrose Esters of Fatty Acids.

## PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store in a cool, dry place free from direct sunlight.
- (2) Keep all medication out of the reach of children.
- (3) When you take this medicine in divided dose, fold over the opened edge of the packet and store it.  
The remaining medicine should be taken within 2 days.
- (4) Since the medicine contains crude drugs, the granules color may vary for individual products, however this has no influence on the efficacy of this medicine.
- (5) Do not take this medicine after the expiration date.

*Production of this product has been approved for domestic sales in accordance with the Pharmaceutical Affairs Law of Japan.*



**Tsumura&Co.** 2-17-11, Akasaka, Minato-ku, Tokyo107-8521, JAPAN  
<https://www.tsumura.co.jp/>